

# Quinone Binding Sites of Membrane Proteins as Targets for Inhibitors\*

Peter R. Rich

Glynn Research Foundation, Glynn, Bodmin, Cornwall PL30 4AU, UK


(Received 17 July 1995; revised version received 15 December 1995; accepted 22 January 1996)

**Abstract:** Quinones are a central component of most photosynthetic and respiratory electron transfer chains. The proteins that reduce and oxidise these quinones have quinone binding sites—Q sites—that are good targets for pesticides. This paper reviews the diversity of these sites, their possible structure, and the types of compounds that act upon them.

**Key words:** biological electron transfer, Q sites, quinones, inhibitors, respiration, photosynthesis

## 1 INTRODUCTION

A common feature of most biological respiratory and photosynthetic electron-transfer chains is that they utilise at some stage the reduction and oxidation of quinones by redox proteins.<sup>1</sup> The three most commonly occurring quinones are shown in Fig. 1. Quinones are often one of the essential elements of the energy coupling process; by using the mobility of quinones between quinone-reactive sites (Q sites) arranged on proteins in a transmembrane manner, the coupling of electron



**Fig. 1.** The structures of biological quinones. Ubiquinone is a component of mitochondrial and many bacterial respiratory chains, whereas plastoquinone is found in higher plant and algal photosynthesis. Menaquinone can be found in place of, or in addition to, ubiquinone in bacterial systems. All forms are extremely hydrophobic because of the long poly-isoprenyl side chain:  $n$  is generally between 8 and 10 for ubiquinone and menaquinone and 9 for plastoquinone.

\* Based on a paper presented at the Symposium 'Interference with the Structure and Function of Biological Membranes,' organised by the Biochemical & Biophysical Panel of the SCI Pesticides Group and held at 14/15 Belgrave Square, London on 23 May 1995.

transfer and proton translocation is achieved as part of a classical redox loop mechanism first described by Mitchell.<sup>2</sup>

Because of the central role of quinones in electron-transfer chains, many different Q sites are known. For many, rapid movement of the quinone on and off the site occurs and the binding constant is relatively weak. This dynamic association is required if the quinone is to act to couple electron and proton transfer, and it provides the possibility of inhibition of the transiently unoccupied site by a quinone analogue or other molecule. Other Q sites involve relatively immobile quinones, and displacement by an antagonist is more difficult. In this case the quinone remains bound during its redox cycle and acts simply as a component of the intra-protein electron-transfer chain.

Both synthetic and natural antagonists are known for a number of these sites in photosynthesis and respiration. All antagonists are relatively hydrophobic and considerable specificity for individual Q sites is often found. Differences in sensitivity of the same site in different species are known. Even a single amino acid change, which does not affect the catalytic activity of the enzyme, can cause a dramatic change in inhibitor sensitivity, and this has allowed the use of genetic methods to probe Q sites by analysis of inhibitor-resistant mutants.<sup>3,4</sup>

The purpose of this article is to review some of the wide range of Q sites that are already known and the types of inhibitor that act on them.

## 2 THE STRUCTURE OF Q SITES

The binding sites of many cofactors or prosthetic groups often involve a characteristic protein fold whose primary sequence has a characteristic pattern for all such sites. The recognition of these protein sequence motifs has been of great value in protein structure prediction. To date, however, the high resolution structures of only two such Q sites in a membrane-bound protein are known, namely the  $Q_A$  and  $Q_B$  sites of the bacterial reaction centre<sup>5,6</sup> and these pockets are formed predominantly not by a single protein fold, but rather by a number of residues in the M ( $Q_A$  site) and L ( $Q_B$  site) subunits that are quite separated in their primary sequences. Hence, there is as yet no general sequence motif for a Q site for use as a tool to search for Q sites in proteins of unknown tertiary structure. Whether such a motif may become evident remains to be established when further Q site structures are elucidated.

Despite the lack of a clear motif, a number of common physical features of Q sites are likely. All such sites are relatively hydrophobic and are accessed by the quinone from the membrane phase, as necessitated by the hydrophobic nature of the quinone substrate. It seems likely that the predominant factor for binding into the site is the provision of groups at the correct distance so that hydrogen bonds can be formed with the quinone/quinol carbonyl/hydroxyl groups. For those sites (the majority) which interconvert the quinone, Q, and the protonated quinol,  $QH_2$ , access of the sites to an aqueous phase is required so that associated protonation reactions can occur. This could be direct or *via* a hydrophilic channel through the protein structure. In the case of the  $Q_B$  site, protonatable groups close to the quinone (GluL212 and AspL213), and polar routes of contact with the aqueous surface, can be recognised.<sup>7</sup> Lack of such access to protons radically changes the properties of the quinone in the site, such that it generally can only cycle between the oxidised, Q, and semi-quinone,  $Q^{\cdot-}$ , forms.

A comparison of the structures around the  $Q_A$  and  $Q_B$  pockets<sup>8</sup> reveals several similarities ( $Q_A$  and  $Q_B$  residues given in brackets):

(i) a histidine group (HisM219 and HisL190) which is ligated to the common iron atom and which is likely to hydrogen bond to one of the quinone carbonyl groups. At least in the  $Q_A$  site, further hydrogen bonding of this carbonyl to the backbone nitrogen of AlaM260 (GlyL225 in  $Q_B$ ) occurs;

(ii) hydrogen bonds between the protein and the second quinone carbonyl group. Originally, these hydrogen bonds were thought to arise from the uncharged polar groups, ThrM222 and SerL223. However, a more recent refinement of the structure of the  $Q_B$  site indicates that hydrogen bonding may instead be to the backbone peptide nitrogens of IleL224 and GlyL225;<sup>9</sup>

(iii) an aromatic residue (TrpM252 and PheL216) which is particularly close to the quinone ring in the  $Q_A$  site, but further away in the  $Q_B$  sites. An aliphatic residue (ValM226 and IleL229) occupies the opposite side of the quinone ring.<sup>10</sup>


The majority of the differences between the physical properties of quinone when bound in the  $Q_A$  and  $Q_B$  sites are likely to arise from the greater hydrophobicity, the more restricted access for quinone exchange, and the lack of protonatable groups of the  $Q_A$  site.

Since the M and L subunits are evolutionarily related, it is impossible at this stage to speculate with confidence on how many of the common features will be shared by other Q sites. However, it seems likely that some new structures which contain Q sites will be elucidated in the near future, and this may allow identification of a motif for recognition of other quinone binding sites. Recently, the crystal structures of mammalian<sup>11</sup> and bacterial forms of a soluble NAD(P)H quinone oxidoreductase have been determined. The structure of the mammalian enzyme with bound duroquinone has also been elucidated and this provides a third Q site structure, albeit of a soluble enzyme. Interestingly, this structure has features in common with the  $Q_A$  and  $Q_B$  sites in terms of hydrogen bonding (in this case of one quinone carbonyl to two tyrosine residues), interaction of the quinone ring with a coplanar ring structure (in this case the isoalloxazine ring of the flavin), and inhibition by Q site inhibitors.<sup>12</sup>

## 3 TYPES OF INHIBITOR

Inhibitors of Q sites are extremely diverse. Many are relatively simple (Fig. 2), with a clear resemblance to the structure of the natural quinones in terms of positions of hydrogen bonding groups and hydrophobic character. Some are redox inactive, at least in their mode of action (although unrelated electrochemical activity at extreme potentials is not uncommon). Such molecules have been discovered from natural, e.g. aurachins,<sup>13</sup> piericidin A<sup>14</sup> and synthetic, e.g. triazines and DCMU (3-(3',4'-dichlorophenyl)-1,1'-dimethylurea), sources. It seems almost inevitable that these compounds bind in place of the quinone in the Q site. In some instances, structured water molecules may also be required to form the stable, inhibitory complex, as has shown to be the case for the binding of atrazine (a triazine) in the  $Q_B$  pocket of the bacterial reaction centre.<sup>15</sup>


Some of these simple molecules are themselves quinones or other redox active compounds. Although they presumably bind to the Q site in the same manner as substrate quinone, their physical properties are such that catalytic turnover cannot be completed. Many synthetic quinones are known in this category e.g. DBMIB (2,5-dibromo-3-methyl-6-isopropyl-*p*-benzoquinone) and various hydroxy-naphthoquinones. In some instances, a partial redox reaction can still occur. For



**Fig. 2.** Structures of some simple Q site inhibitors. Further information on these and related derivatives can be found as follows: DBMIB,<sup>89</sup> alkyl-HQNO;<sup>79,90</sup> flavones;<sup>68</sup> hydroxypyridines;<sup>89</sup> alkylgallates;<sup>33,34</sup> UHDBT;<sup>91</sup> alkyl-2-hydroxynaphthoquinones;<sup>39</sup> acridones;<sup>64,92</sup> aurachins;<sup>13,93</sup> piericidin A;<sup>14</sup> triazines and phenylureas.<sup>37,89</sup>

example, DBMIB is usually converted to its reduced state, DBMIBH<sub>2</sub>, when added to a biological system such as chloroplasts, and is not an inhibitor of the Q sites of chloroplast cytochrome *bc*<sub>1</sub> complex in this form. However, oxidation at one of the Q sites converts it into a bound form which is an extremely potent inhibitor.<sup>16</sup> In other instances, redox active chemicals may not bind strongly to a Q site at all, but show biological toxicity because they are biochemically reduced at the site and then autoxidise to produce toxic species such as superoxide anions.

Other Q site inhibitors have a more complex structure, and regions capable of binding in place of the substrate quinone are not so clear. Although direct binding to the Q site may still occur, significant interaction with the protein structure surrounding the Q site seems likely. Inhibition again probably occurs by blocking the binding of the natural quinone, or possibly its access route, to the site, although in some instances a more allosteric effect on the Q site has been postulated. The more complex molecules have been discovered initially from natural sources, e.g. antimycin, myxothiazol, rotenone and acetogenins such as bullatacin (Fig. 3 and Table 1). For two cases that have been tested,<sup>9,17</sup> the detailed molecular stereochemistry (Fig. 4) has been shown to be critical for activity. However, the full complexity of some natural compounds is not always essential. In the case of stigmatellin (Fig. 3), for example, the complicated tail can be replaced by a simple *n*-alkyl one of appropriate partition coefficient, and the crystal



**Fig. 3.** Structures of some mitochondrial Q site inhibitors. References to further information on these compounds are given in Table 1.


**TABLE 1**  
Q Sites and Inhibitors in Mitochondria

Enzyme	Likely number of Q sites	Examples of tight, <sup>a</sup> relatively specific <sup>b</sup> inhibitors	Other inhibitors or relevant literature
Complex I <sup>c</sup>	One or two <sup>56,57</sup>	Rotenone <sup>14</sup> Piericidin A <sup>14</sup> Acetogenins <sup>58,59</sup> Fenazaquin <sup>27</sup> Capsaicin derivatives <sup>60</sup>	MPP <sup>+</sup> 61–63 Others. <sup>9,26,27,57,64</sup>
NDH-2 <sup>65</sup>	One	None	Refs 65, 66
NADH <sub>ext</sub> dehydrogenase <sup>67</sup>	One	None	Flavones <sup>68</sup>
Succinate dehydrogenase	Two <sup>69</sup>	Carboxins <sup>70–72</sup>	TTFA <sup>71</sup>
Electron-transferring flavoprotein <sup>73,74</sup>	One	None	
D-glycerol-3-phosphate dehydrogenase <sup>74</sup>	One	None	
bc <sub>1</sub> complex	Two <sup>35</sup> or three <sup>75</sup>	Antimycin A <sup>76</sup> Myxothiazol <sup>77</sup> Methoxyacrylate derivatives including oudemansins and strobilurins <sup>78</sup> Funiculosin <sup>79</sup> Stigmatellins <sup>80</sup>	Refs 81, 82
Ubiquinol oxidase <sup>83</sup>	At least one	Alkyl gallates <sup>34</sup>	Refs 30–32

<sup>a</sup> Tight binding is arbitrarily defined as an observed dissociation constant of  $10^{-6}$  M or less in at least one species. However, dramatic variation between species can occur and such dissociation constants are in any case dependent on assay conditions and type of enzyme preparation.

<sup>b</sup> Showing a reasonable specificity for one Q site within the mitochondrial electron transfer chain, although cross-reactivity with Q sites in other electron-transfer systems occurs in some cases.

<sup>c</sup> NADH dehydrogenases are categorised as complex I (or NDH-1), NDH-2 (by analogy with categorisation of a second bacterial enzyme<sup>65</sup> and a third enzyme, NADH<sub>ext</sub> dehydrogenase, found in plants and fungi which oxidises cytosolic NADH and presumed to be a distinct enzyme type.



**Fig. 4.** Conformations of some complex Q site inhibitors. The structures of two natural Q site inhibitors, rotenone and (+)-antimycin A<sub>1</sub>, have been computed using the HYPERCHEM energy minimisation software. The calculated rotenone structure agrees well with that determined crystallographically.<sup>94</sup>

structure of a reaction centre-stigmatellin complex clearly shows that only the headgroup has specific interactions with the protein, to which it binds in a quinone-like manner.<sup>15</sup>

Regardless of complexity of structure, some inhibitors are tight and immobile, staying attached to a single site for many seconds, whereas others are tight but rapidly mobile on a millisecond timescale between identical sites in the same membrane.<sup>18</sup> The degree of inhibitor mobility is not directly correlated with quinone exchangeability in the site, and the factors which control the rate constants of inhibitor exchange are not understood.

## 4 SOME Q SITES AND THEIR INHIBITORS

### 4.1 All mitochondrial respiratory chains


The majority of mitochondrial respiratory chains possess a common basic set of components, and these

have perhaps been most studied in the mammalian mitochondrion. The pool of ubiquinone connects dehydrogenases such as NADH dehydrogenase (complex I) and succinate dehydrogenase (complex II) to the cytochrome  $bc_1$  complex (complex III) and all three of these enzymes and their Q site reactivities have been studied in detail. Although a number of powerful inhibitors are already known for the majority of their known Q sites (Table 1 and Fig. 3), specificity for the enzyme in particular organisms, e.g. for fungal or insect forms of the enzymes, remains a possibility. For example, specificity of benzimidazole-*N*-sulfonamides for the  $bc_1$  complex of *Pythium aphanidermtum* has been found.<sup>19</sup> In some instances, insensitivity of a species to a compound which inhibits the same enzyme in closely related organisms has been shown, e.g. an insensitivity to the  $bc$  complex in sea urchins to the classical inhibitors myxothiazol and strobilurin<sup>20</sup> or in fish to funiculosin.<sup>21</sup> Commercial applications of some compounds that act on these Q sites have been described.<sup>22–27</sup>

## 4.2 Plant, yeast and fungal mitochondria

Mitochondria from yeasts, fungi and higher plants can contain additional enzymes that are not found in mammalian systems (Fig. 5). In particular, a very active externally facing, inner membrane NADH dehydrogenase is often present. In addition, a second NADH dehydrogenase (termed NDH-2 in order to distinguish it from complex I) for oxidation of internally generated NADH is present which is insensitive to complex I inhibitors.<sup>28</sup> Although a variety of rather weak Q site inhibitors are known for these additional NADH dehydrogenases, no tight and specific inhibitors are yet available.

In addition to the extra dehydrogenases, mitochondria from these sources often possess a second 'alterna-



**Fig. 5.** Q sites of proteins of mitochondrial respiratory chains. Q sites are indicated by shaded semicircles. Arrows indicate normal direction of electron flow. A question mark indicates uncertainty on the existence of the site. Components are not universally present in mitochondria from different species. UQ = ubiquinone; c = cytochrome c.


tive' oxidase which oxidises ubiquinol directly.<sup>29</sup> A variety of rather weak or non-specific inhibitors have been known for some time.<sup>30</sup> One of those most commonly used at present is *n*-propyl gallate, although concentrations in excess of  $10^{-5}$  M are required for maximum inhibition.<sup>31</sup> Substitution of the *n*-propyl group for an octyl or dodecyl group increased the potency of inhibition of an analogous trypanosomal enzyme<sup>32,33</sup> and we found recently that the same change increased substantially the potency of inhibition of the plant enzyme, so that the inhibitor became stoichiometric in some conditions.<sup>34</sup>

## 4.3 Higher plant and algal chloroplasts

The higher plant photosynthetic electron-transfer chain consists of two photosystems, connected in series by plastoquinone, a cytochrome *bf* complex (homologous to the mitochondrial  $bc_1$  complex) and plastocyanin. More than ten Q sites have been proposed,<sup>35</sup> but not all have withstood the test of time. Six of them are now well-documented (Fig. 6). However, a number of other functions recently recognised in thylakoids imply that other novel Q sites are likely to be found in the future.

Two of the sites are on photosystem II and are functionally similar to the  $Q_A$  and  $Q_B$  sites in the bacterial reaction centre. Indeed, the sequence homology between photosystem II and the bacterial reaction centre also suggests that they are similar, and structural models have been proposed on this basis, see Ref. 36. Because of its inaccessibility and the lack of exchangeability of the plastoquinone when bound, the  $Q_A$  site is not a good target for inhibitors. In contrast, a wide range of inhibitors can bind to the  $Q_B$  site and prevent reaction with plastoquinone.<sup>36–40</sup> Indeed, many commercially important herbicides, such as the triazines and phenylureas (Fig. 2), act on this site.

Two further Q sites, termed the  $Q_o$  and the  $Q_i$  sites, occur on the cytochrome *bf* complex. This enzyme is homologous to the mitochondrial and bacterial *bc* complexes and it is likely that the structures and function of the two Q sites are similar. However, the majority of the



**Fig. 6.** Q sites in higher plant thylakoids. Conventions are as for Fig. 5. PQ = plastoquinone; PC = plastocyanin; Fd = ferredoxin; FQR = a putative ferredoxin-plastoquinone oxidoreductase.

very tight, specific, inhibitors of the Q sites of the *bc* complexes do not act on the chloroplast enzyme. Of the 'classical' mitochondrial Q<sub>o</sub> site inhibitors, only stigma-tellin (Fig. 4) acts in the same way on the cytochrome *bf* complex<sup>1</sup> and recently we have found that its potency can be improved by exchange of the natural sidechain for a simpler C<sub>13</sub> *n*-alkyl one.<sup>41</sup> None of the classical *bc* complex inhibitors acts well on the Q<sub>i</sub> site of the cytochrome *bf* complex. The first compounds shown to act on this site were HQNO and NQHO (2-*n*-heptyl-4-hydroxyquinoline-*N*-oxide and 2-*n*-nonyl-4-hydroxyquinoline-*N*-oxide),<sup>42</sup> and we found improved specificity with MOA-stilbene (*E*- $\beta$ -methoxyacrylate-stilbene or (*E,E*)-methyl 3-methoxy-2-(styrylphenyl)propenoate)<sup>43</sup> and, more recently, with a compound that we have termed FIQ (5-(3',4'-dimethoxybenzyl)-2-(1'-methyl-2'-phenylthioethyl)-2,3,5,6,7,8-hexahydrofuro[2,3-*g*]isoquinoline).<sup>41</sup> However, we still have not achieved the potency and specificity seen with corresponding inhibitors of the Q<sub>i</sub> site of the *bc* complexes. This, in combination with a number of other differences shown by the *bf* complexes, has even led some workers to consider that the chloroplast enzyme might work in a quite different way to the *bc* complexes, although we have argued that this view is not yet justified.<sup>44</sup> In any case, there is still a need to identify more powerful inhibitors of this site.

A further Q site appears within the photosystem I complex. The quinone, termed A<sub>1</sub>, is a naphthoquinone, presumably tightly bound in a rather aprotic and apolar environment, and acting as a simple electron-transfer component between chlorophyll and the iron sulphur centres by cycling between its oxidised and semiquinone forms.<sup>45</sup> Although such a site might be presumed to be difficult to inhibit competitively, such studies have been reported,<sup>46,47</sup> and this site remains a novel target.

It is known that photosynthetic ATP production can occur by electron transfer around photosystem I without the involvement of photosystem II. This process of cyclic photophosphorylation may be the only one that occurs in some types of plant cell<sup>48</sup> but the degree to which it occurs in more conventional thylakoids remains uncertain. It is known to require ferredoxin, which accepts an electron from photosystem I and passes it back to plastoquinone *via* a ferredoxin-plastoquinone oxidoreductase termed FQR.<sup>49</sup> Although this FQR protein has not yet been characterized, it does appear to have a unique Q site which interacts with plastoquinone and it is this site which is likely to account for the sensitivity of the process to antimycin A and other likely Q site inhibitors.<sup>50,51</sup> Again, a more thorough investigation of other possible inhibitors of this Q site is warranted.

Besides these sites, there are a number of more enigmatic Q sites that are likely to be operative in the more intact *in-vivo* situation. For example, a number of addi-

tional sites in intact cells may be present which control a slow reduction and reoxidation of the plastoquinone pool which can occur in the dark in response to other physiological changes of the cell. These processes have collectively been termed 'chlororespiration' and have been suggested to arise from specific thylakoid forms of NADH dehydrogenase and plastoquinol oxidase.<sup>52</sup> However, direct evidence for such enzymes in higher plant or algal thylakoids is poor and the molecular nature of dark plastoquinone reactions has yet to be established. A further example is provided by a major medium-term control mechanism of protein phosphorylation/dephosphorylation, which controls the balance of photosystem activities and responds to the redox state of the plastoquinone pool.<sup>53</sup> This implies that the proteins involved in sensing redox state changes are likely to have sites specific for the Q or QH<sub>2</sub> forms of plastoquinone. The identification of specific inhibitors for any of these sites would provide a major tool for understanding their molecular origin and might in turn provide new target sites for herbicides.

#### 4.4 Bacterial systems

Since the mitochondrial and thylakoid systems evolved from bacterial origins, it is not surprising that homologues of the majority of enzymes described above can be found in the electron-transfer chains of various bacterial species. Generally, inhibitors that act on the eukaryotic enzymes are also effective on their bacterial counterparts. In addition, however, there is a range of more specialised enzymes in different bacteria for the reduction or oxidation of the quinone by diverse metabolic substrates, and more are likely to be discovered. Table 2 summarises some of these bacterial enzymes. Even for a particular type of enzyme, great diversity of

TABLE 2  
Some Additional Bacterial Enzymes Which May Have Q Sites

Enzyme	Donor or acceptor	References
<i>Quinone-Reducing<sup>a</sup></i>		
Hydrogenase	Hydrogen	74
NDH-2	NADH	65
Na <sup>+</sup> -NQR	NADH	84
Formate dehydrogenase	Formate	85
Glucose dehydrogenase	Glucose	86
<i>Quinol-Oxidising<sup>a</sup></i>		
Cytochrome <i>bo</i>	Oxygen	87
Cytochrome <i>bd</i>	Oxygen	87
Nitrate reductase	Nitrate	74
DMSO reductase	Dimethyl sulphoxide	74
TMAO reductase	Trimethylamine- <i>N</i> -oxide	74
Fumarate reductase	Fumarate	88

<sup>a</sup> The quinone system in bacteria may be ubiquinone, menaquinone or both.

detail is found between homologues in different bacterial species (e.g. the variants of the oxidase superfamily<sup>54</sup>). Recently, we screened a range of compounds as possible inhibitors of the Q sites of the two principal bacterial quinol oxidases, cytochromes *bo* and *bd*, and identified a number of compounds that are powerful inhibitors of these sites.<sup>55</sup> However, inhibitors of the Q sites of many of the other bacterial enzymes have yet to be identified.

## 5 CONCLUSIONS

The Q sites form a wide and diverse class of target sites for specific inhibitors and it has yet to be established whether they have a common primary sequence motif. Although they are generally rather simple and non-specific for the types of quinone that they can recognise, great specificity for individual sites is often exhibited by inhibitors. The search for new inhibitors offers exciting prospects both for aiding the elucidation of the molecular properties of the enzymes and for providing new types of commercially viable pesticides.

## ACKNOWLEDGEMENTS

Part of this work is funded by a research grant (Ref. 101/PO2632) from the BBSRC.

## REFERENCES

- Rich, P. R., Electron and proton transfers through quinones and cytochrome *bc* complexes. *Biochim. Biophys. Acta*, **768** (1984) 53–79.
- Mitchell, P., Coupling of phosphorylation to electron and proton transfer by a chemi-osmotic type of mechanism. *Nature (London)*, **191** (1961) 144–8.
- Degli Esposti, M., de Vries, S., Crimi, M., Ghelli, A., Patarnello, T. & Meyer, A., Mitochondrial cytochrome *b*: Evolution and structure of the protein. *Biochim. Biophys. Acta*, **1143** (1993) 243–71.
- Colson, A.-M., Random mutant generation and its utility in uncovering structural and functional features of cytochrome *b* in *Saccharomyces cerevisiae*. *J. Bioenerg. Biomemb.*, **25** (1993) 211–20.
- Deisenhofer, J., Epp, O., Miki, K., Huber, R. & Michel, H., Structure of the protein subunits in the photosynthetic reaction centre of *Rhodospseudomonas viridis* at 3 Å resolution. *Nature (London)*, **318** (1985) 618–24.
- Feher, G., Allen, J. P., Okamura, M. Y. & Rees, D. C., Structure and function of bacterial photosynthetic reaction centres. *Nature (London)*, **339** (1989) 111–6.
- Okamura, M. Y. & Feher, G., Proton transfer in reaction centers from photosynthetic bacteria. *Ann. Rev. Biochem.*, **61** (1992) 861–96.
- Allen, J. P., Feher, G., Yeates, T. O., Komiya, H. & Rees, D. C., Structure of the reaction center from *Rhodobacter sphaeroides* R-26: Protein-cofactor (quinones and Fe<sup>2+</sup>) interactions. *Proc. Natl Acad. Sci. USA*, **85** (1988) 8487–91.
- Ueno, H., Miyoshi, H., Ebisui, K. & Iwamura, H., Comparison of the inhibitory action of natural rotenone and its stereoisomers with various NADH-ubiquinone reductases. *Eur. J. Biochem.*, **225** (1994) 411–7.
- Robertson, D. E., Daldal, F. & Dutton, P. L., Mutants of ubiquinol cyt c2 oxidoreductase resistant to Qo site inhibitors: consequences for ubiquinone and ubiquinol affinity and catalysis. *Biochemistry*, **29** (1990) 11249–60.
- Li, R. B., Bianchet, M. A., Talalay, P. & Amzel, L. M., The three-dimensional structure of NAD(P)H-quinone reductase, a flavoprotein involved in cancer chemoprotection and chemotherapy: Mechanism of the two-electron reduction. *Proc. Natl Acad. Sci. USA*, **92** (1995) 8846–50.
- Chen, S., Hwang, J. & Deng, P. S. K., Inhibition of NAD(P)H: quinone acceptor oxidoreductase by flavones: A structure-activity study. *Arch. Biochem. Biophys.*, **302** (1993) 72–7.
- Kunze, B., Höfle, G. & Reichenbach, H., The aurachins, new quinoline antibiotics from myxobacteria: production, physicochemical and biological properties. *J. Antibiotics*, **40** (1987) 258–65.
- Singer, T. P., Mitochondrial electron transport inhibitors. *Meth. Enzymol.*, **55** (1979) 454–65.
- Lancaster, C. R. D. & Michel, H., New insights into the X-ray structure of the reaction center from *Rhodospseudomonas viridis*. In *Reaction Centers of Photosynthetic Bacteria. Structure and Dynamics. International Workshop Feldafing III*, ed. M. E. Michel-Beyerle, M. E. Springer-Verlag, Berlin, 1995.
- Rich, P. R., Madgwick, S. A. & Moss, D. A., The interactions of duroquinol, DBMIB and NQNO with the chloroplast cytochrome *bf* complex. *Biochim. Biophys. Acta*, **1058** (1991) 312–28.
- Miyoshi, H., Kondo, H., Oritani, T., Saitoh, I. & Iwamura, H., Inhibition of electron transport of rat liver mitochondria by unnatural (–)-antimycin A<sub>3</sub>. *FEBS Lett.*, **292** (1991) 61–3.
- Bechmann, G., Weiss, H. & Rich, P. R., Non-linear inhibition curves for tight-binding inhibitors of dimeric ubiquinol-cytochrome *c* oxidoreductases. Evidence for rapid inhibitor mobility. *Eur. J. Biochem.*, **208** (1992) 315–25.
- Pillonel, C., Interaction of benzimidazole-*N*-sulfonamides with the cytochrome *bc*<sub>1</sub> complex of *Pythium aphididermatum*. *Pestic. Sci.*, **43** (1995) 107–13.
- Degli Esposti, M., Ghelli, A., Butler, G., Roberti, M., Mustich, A. & Cantatore, P., The cytochrome *b* of the sea urchin *Paracentrotus lividus* is naturally resistant to myxothiazol and mucidin. *FEBS Lett.*, **263** (1990) 245–7.
- Degli Esposti, M., Ghelli, A., Crimi, M., Baracca, A., Solaini, G., Tron, T. & Meyer, A., Cytochrome *b* of fish mitochondria is strongly resistant to funiculosin, a powerful inhibitor of respiration. *Arch. Biochem. Biophys.*, **295** (1992) 198–204.
- Mathre, D. E., Mode of action of oxathiin fungicides. I. Effects of carboxin and oxycarboxin on the general metabolism of several basidiomycetes. *Phytopathology*, **60** (1970) 671–6.
- Ammermann, E., Lorenz, G., Schelberger, K., Wenderoth, B., Sauter, H. & Rentzea, C., BAS 490 F—a broad-spectrum fungicide with a new mode of action. *Brighton Crop Prot. Conf.* (1992) 403–10.
- Beautement, K., Clough, J. M., de Fraine, P. J. & Godfrey, C. R. A., Fungicidal *b*-methoxyacrylates: from natural products to novel synthetic agricultural fungicides. *Pestic. Sci.*, **31** (1991) 499–519.
- Clough, J. M. & Godfrey, C. R. A., Growing hopes. *Chemistry in Britain*, **31** (1995) 466–9.
- Jewess, P. J., Insecticides and acaricides which act at the rotenone-binding site of mitochondrial

- NADH:ubiquinone oxidoreductase; Competitive displacement studies using a  $^3\text{H}$ -labelled rotenone analogue. *Biochem. Soc. Trans.*, **22** (1994) 247–51.
27. Hollingworth, R. M., Ahammadsahib, K. I., Gadelhak, G. & McLaughlin, J. L., New inhibitors of Complex I of the mitochondrial electron transport chain with activity as pesticides. *Biochem. Soc. Trans.*, **22** (1994) 230–3.
  28. Menz, R. I., Griffith, M., Day, D. A. & Wiskich, J. T., Matrix NADH dehydrogenases of plant mitochondria and sites of quinone reduction by complex I. *Eur. J. Biochem.*, **208** (1992) 481–5.
  29. Bonner, W. D., Jr, Clarke, S. D. & Rich, P. R., Partial purification and characterisation of the quinol oxidase activity of *Arum maculatum* mitochondria. *Plant Physiol.*, **80** (1986) 838–42.
  30. Schonbaum, G. R., Bonner, W. D., Jr, Storey, B. T. & Bahr, J. T., Specific inhibition of the cyanide-insensitive respiratory pathway in plant mitochondria by hydroxamic acids. *Plant Physiol.*, **47** (1971) 124–8.
  31. Siedow, J. N. & Girvin, M. E., Alternative respiratory pathway: Its role in seed respiration and its inhibition by propyl gallate. *Plant Physiol.*, **65** (1980) 669–74.
  32. Grady, R. W., Bienen, E. J., Dieck, H. A., Saric, M. & Clarkson, A. B., Jr, *N-n*-Alkyl-3,4-dihydroxybenzamides as inhibitors of the trypanosome alternative oxidase: activity *in vitro* and *in vivo*. *Antimicrobial Agents and Chemotherapy*, **37** (1993) 1082–5.
  33. Grady, R. W., Bienen, E. J. & Clarkson, A. B., Jr, Esters of 3,4-dihydroxybenzoic acid, highly effective inhibitors of the *sn*-glycerol-3-phosphate oxidase of *Trypanosoma brucei brucei*. *Mol. Biochem. Parasitol.*, **21** (1986) 55–63.
  34. Hoefnagel, M. H. N., Wiskich, J. T., Madgwick, S. A., Patterson, Z., Oettmeier, W. & Rich, P. R., New inhibitors of the ubiquinol oxidase of higher plants. *Eur. J. Biochem.*, **233** (1995) 531–7.
  35. Rich, P. R. & Moss, D. A., The reactions of quinones in higher plant photosynthesis. In *The Light Reactions*, ed. J. Barber. Elsevier Science Publishers BV, Amsterdam, 1987, pp. 421–445.
  36. Tietjen, K. G., Kluth, J. F., Andree, R., Haug, M., Lindig, M., Müller, K. H., Wroblowsky, H. J. & Trebst, A., The herbicide binding niche of photosystem II—a model. *Pestic. Sci.*, **31** (1991) 65–72.
  37. Fedtke, C., *Biochemistry and Physiology of Herbicide Action*. Springer-Verlag, Berlin, 1992.
  38. Sobolev, V. & Edelman, M., Modeling the quinone-B binding site of the photosystem-II reaction center using notions of complementarity and contact-surface between atoms. *Proteins*, **21** (1995) 214–25.
  39. Draber, W., Pittel, B. & Trebst, A., Modeling of photosystem II inhibitors of the herbicide-binding protein. In *Probing Bioactive Mechanisms*. ACS Symposium Series No. 413, ed. P. S. Magee, D. R. Henry & J. H. Block, 1991, pp. 215–228.
  40. Warncke, K., Gunner, M. R., Braun, B. S., Gu, L., Yu, C.-A., Bruce, J. M. & Dutton, P. L., Influence of hydrocarbon tail structure on quinone binding and electron-transfer performance of the  $\text{Q}_\text{A}$  and  $\text{Q}_\text{B}$  sites of the photosynthetic reaction center protein. *Biochemistry*, **33** (1994) 7830–41.
  41. Ondarroa, M., Madgwick, S. A., Wiggins, T. E., Whiting, D. A. & Rich, P. R., Improved inhibitors of the cytochrome *bf* complex of higher plants. *Abstracts of the Eighth European Bioenergetics Conference, Valencia, Spain* (1994), p. 59.
  42. Jones, R. W. & Whitmarsh, J., Inhibition of electron transfer and the electrogenic reaction in the cytochrome *b/f* complex by 2-*n*-nonyl-4-hydroxyquinoline *N*-oxide (NQNO) and 2,5-dibromo-3-methyl-6-isopropyl-*p*-benzoquinone (DBMIB). *Biochim. Biophys. Acta*, **933** (1988) 258–68.
  43. Rich, P. R., Madgwick, S. A., Brown, S., von Jagow, G. & Brandt, U., MOA-stilbene: A new tool for investigation of the reactions of the chloroplast cytochrome *bf* complex. *Photosynth. Res.*, **34** (1992) 465–77.
  44. Rich, P. R., A critical examination of the supposed variable proton stoichiometry of the chloroplast cytochrome *bf* complex. *Biochim. Biophys. Acta*, **932** (1988) 33–42.
  45. Snyder, S. W., Rustandi, R. R., Biggins, J., Norris, J. R. & Thurnauer, M. C., Direct assignment of vitamin  $\text{K}_1$  as the secondary acceptor  $\text{A}_1$  in photosystem I. *Proc. Natl Acad. Sci. USA*, **88** (1991) 9895–6.
  46. Itoh, S. & Iwaki, M., Full replacement of the function of the secondary electron acceptor phyloquinone( = vitamin  $\text{K}_1$ ) by non-quinone carbonyl compounds in green plant photosystem I photosynthetic reaction centers. *Biochemistry*, **30** (1991) 5340–6.
  47. Iwaki, M. & Itoh, S., Structure of the phyloquinone-binding ( $\text{Q}_{\text{phl}}$ ) site in green plant photosystem I reaction centers: The affinity of quinones and quinonoid compounds for the  $\text{Q}_{\text{phl}}$  site. *Biochemistry*, **30** (1991) 5347–52.
  48. Bendall, D. S. & Manasse, R. S., Cyclic phosphorylation and electron transport. *Biochim. Biophys. Acta*, **1229** (1995) 23–38.
  49. Cleland, R. E. & Bendall, D. S., Photosystem I cyclic electron transport: Measurement of ferredoxin-plastoquinone reductase activity. *Photosynth. Res.*, **34** (1992) 409–18.
  50. Trebst, A. & Wietoska, H., On an inhibitor of ferredoxin-catalyzed cyclic photophosphorylation in thylakoid membranes. *Botanica Acta*, **102** (1989) 302–5.
  51. Patel, P. K., Bendall, D. S. & Ridley, S. M., Site of action of a halogenated 4-hydroxypyridine on ferredoxin-catalysed cyclic photophosphorylation. *FEBS Lett.*, **206** (1986) 249–52.
  52. Bennoun, P., Evidence for a respiratory chain in the chloroplast. *Proc. Natl Acad. Sci. USA*, **79** (1982) 4352–6.
  53. Allen, J. F., Protein phosphorylation in regulation of photosynthesis. *Biochim. Biophys. Acta*, **1098** (1992) 275–335.
  54. Saraste, M., Holm, L., Lemieux, L., Lübben, M. & van der Oost, J., The happy family of cytochrome oxidases. *Biochem. Soc. Trans.*, **19** (1991) 608–12.
  55. Meunier, B., Madgwick, S. A., Reil, E., Oettmeier, W. & Rich, P. R., New inhibitors of the quinol oxidation sites of bacterial cytochromes *bo* and *bd*. *Biochemistry*, **34** (1995) 1076–83.
  56. Degli Esposti, M. D. & Ghelli, A., The mechanism of proton and electron transport in mitochondrial complex I. *Biochim. Biophys. Acta*, **1187** (1994) 116–20.
  57. Friedrich, T., Van Heek, P., Leif, H., Ohnishi, T., Forche, E., Kunze, B., Jansen, R., Trowitzsch-Kienast, W., Höfle, G., Reichenbach, H. & Weiss, H., Two binding sites of inhibitors in NADH:ubiquinone oxidoreductase (complex I)—Relationship of one site with the ubiquinone-binding site of bacterial glucose:ubiquinone oxidoreductase. *Eur. J. Biochem.*, **219** (1994) 691–8.
  58. Degli Esposti, M., Ghelli, A., Ratta, M., Cortes, D. & Estornell, E., Natural substances (acetogenins) from the family *Annonaceae* are powerful inhibitors of mitochondrial NADH dehydrogenase (Complex I). *Biochem. J.*, **301** (1994) 161–7.
  59. Londershausen, M., Leicht, W., Lieb, F., Moeschler, H. & Weiss, H., Molecular mode of action of annonins. *Pestic. Sci.*, **33** (1991) 427–38.
  60. Satoh, T., Miyoshi, H., Sakamoto, K. & Iwamura, H., Comparison of the inhibitory action of synthetic capsaicin



- analogues with various NADH-ubiquinone oxidoreductases. *Biochim. Biophys. Acta* (1996) *in press*.
61. Ramsay, R. R., Kowal, A. T., Johnson, M. K., Salach, J. I. & Singer, T. P., The inhibition site of MPP<sup>+</sup>, the neurotoxic bioactivation product of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine is near the Q-binding site of NADH dehydrogenase. *Arch. Biochem. Biophys.*, **259** (1987) 645–9.
  62. Murphy, M. P., Krueger, M. J., Sablin, S. O., Ramsay, R. R. & Singer, T. P., Inhibition of Complex I by hydrophobic analogues of *N*-methyl-4-phenylpyridinium (MPP<sup>+</sup>) and the use of an ion-selective electrode to measure their accumulation by mitochondria and electron-transport particles. *Biochem. J.*, **306** (1995) 359–65.
  63. Ramsay, R. R., Krueger, M. J., Youngster, S. K., Gluck, M. R., Casida, J. E. & Singer, T. P., Interaction of 1-methyl-4-phenylpyridinium ion (MPP<sup>+</sup>) and its analogs with the rotenone/piericidin binding site of NADH dehydrogenase. *J. Neurochem.*, **56** (1991) 1184–90.
  64. Oettmeier, W., Masson, K. & Soll, M., The acridones, new inhibitors of mitochondrial NADH : ubiquinone oxidoreductase (complex I). *Biochim. Biophys. Acta*, **1099** (1992) 262–6.
  65. Yagi, T., The bacterial energy-transducing NADH-quinone oxidoreductases. *Biochim. Biophys. Acta*, **1141** (1993) 1–17.
  66. Bourne, R. M. & Rich, P. R., Characterisation of a sodiummotive NADH : ubiquinone oxidoreductase. *Biochem. Soc. Trans.*, **20** (1992) 577–82.
  67. Moller, I. M., Rasmussen, A. G. & Fredlund, K. M., NAD(P)H-ubiquinone oxidoreductases in plant mitochondria. *J. Bioenerg. Biomemb.*, **25** (1993) 377–84.
  68. Ravanel, P., Creuzet, S. & Tissut, M., Inhibitor effect of hydroxyflavones on the exogenous NADH dehydrogenase of plant mitochondrial inner membranes. *Phytochemistry*, **29** (1990) 441–6.
  69. Ohnishi, T., Structure of the succinate-ubiquinone oxidoreductase (complex II). In *Current Topics in Bioenergetics, Volume 15*, Academic Press, New York, 1987, p. 37.
  70. Coles, C. J., Singer, T. P., White, G. A. & Thorn, G. D., Studies on binding of carboxin analogs to succinate dehydrogenase. *J. Biol. Chem.*, **253** (1978) 5573–8.
  71. Mowery, P. C., Steenkamp, D. J., Ackrell, B. A. C., Singer, T. P. & White, G. A., Inhibition of mammalian succinate dehydrogenase by carboxins. *Arch. Biochem. Biophys.*, **178** (1977) 495–506.
  72. Ackrell, B. A. C., Kearney, E. B., Coles, C. J., Singer, T. P., Beinert, H., Wan, Y.-P. & Folkers, K., Kinetics of reoxidation of succinate dehydrogenase. *Arch. Biochem. Biophys.*, **182** (1977) 107–17.
  73. Ramsay, R. R., Steenkamp, D. J. & Husain, M., Reactions of electron-transfer flavoprotein and electron-transfer flavoprotein : ubiquinone oxidoreductase. *Biochem. J.*, **241** (1987) 883–92.
  74. Nicholls, D. G. & Ferguson, S. J., *Bioenergetics 2*. Academic Press, London, 1992.
  75. Ding, H., Robertson, D. E., Daldal, F. & Dutton, P. L., Cytochrome *bc*<sub>1</sub> complex [2Fe-2S] cluster and its interaction with ubiquinone and ubihydroquinone at the Q<sub>o</sub> site: A double-occupancy Q<sub>o</sub> site model. *Biochemistry*, **31** (1992) 3144–58.
  76. Slater, E. C., The mechanism of action of the respiratory inhibitor antimycin. *Biochim. Biophys. Acta*, **301** (1973) 129–54.
  77. Thierbach, G. & Reichenbach, H., Myxothiazol, a new inhibitor of the cytochrome *bc*<sub>1</sub> segment of the respiratory chain. *Biochim. Biophys. Acta*, **638** (1981) 282–9.
  78. Becker, W. F., von Jagow, G., Anke, T. & Steglich, W., Oudemansin, strobilurin A, strobilurin B and myxothiazol: new inhibitors of the *bc*<sub>1</sub> segment of the respiratory chain with an E- $\beta$ -methoxyacrylate system as common structural element. *FEBS Lett.*, **132** (1981) 329–33.
  79. von Jagow, G., & Link, T. A., Use of specific inhibitors on the mitochondrial *bc*<sub>1</sub> complex. In *Methods in Enzymology* ed. S. Fleischer & B. Fleischer, Academic Press, New York, 1986, pp. 253–271.
  80. von Jagow, G. & Ohnishi, T., The chromone inhibitor stigmatellin—binding to the ubiquinol oxidation center at the C-side of the mitochondrial membrane. *FEBS Lett.*, **185** (1985) 311–15.
  81. von Jagow, G. & Link, T. A., Use of specific inhibitors on the mitochondrial *bc*<sub>1</sub> complex. *Meth. Enzymol.*, **126** (1986) 253–71.
  82. Miyoshi, H., Saitoh, I. & Iwamura, H., Quantitative analysis of electron transport inhibition of rat-liver mitochondrial cytochrome *bc*<sub>1</sub> complex by nitrophenols. *Biochim. Biophys. Acta*, **1143** (1993) 23–8.
  83. Moore, A. L. & Siedow, J. N., The regulation and nature of the cyanide-resistant alternative oxidase of plant mitochondria. *Biochim. Biophys. Acta*, **1059** (1991) 121–40.
  84. Rich, P. R., Meunier, B. & Ward, F. B., Predicted structure and possible ionmotive mechanism of the sodium-linked NADH-ubiquinone oxidoreductase of *Vibrio alginolyticus*. *FEBS Lett.*, **375** (1995) 5–10.
  85. Schlindwein, C., Giordano, G., Santini, C. L. & Mandrand, M.-A., Identification and expression of the *Escherichia coli fdhD* and *fdhE* genes, which are involved in the formation of respiratory formate dehydrogenase. *J. Bacteriol.*, **172** (1990) 6112–21.
  86. van Schie, B. J., Hellingwert, K. J., van Dijken, J. P., Elferink, M. G. L., van Dijk, J. M., Kuenen, J. G. & Konings, W. N., Energy transduction by electron transfer via a pyroloquinoline quinone-dependent glucose dehydrogenase in *Escherichia coli*. *J. Bacteriol.*, **163** (1985) 493–9.
  87. Sato-Watanabe, M., Mogi, T., Miyoshi, H., Iwamura, H., Matsushita, K., Adachi, O. & Anraku, Y., Structure-function studies on the ubiquinol oxidation site of the cytochrome *bo* complex from *Escherichia coli* using *p*-benzoquinones and substituted phenols. *J. Biol. Chem.*, **269** (1994) 28899–907.
  88. Cole, S. T., Condon, C., Lemire, B. D. & Weiner, J. H., Molecular biology, biochemistry and bioenergetics of fumarate reductase, a complex membrane-bound iron-sulfur flavoenzyme of *Escherichia coli*. *Biochim. Biophys. Acta*, **811** (1985) 381–403.
  89. Trebst, A., Inhibitors of electron flow: tools for the functional and structural localization of carriers and energy conservation sites. In *Methods in Enzymology, Volume 69c*, ed. S. P. Colowick & N. O. Kaplan. Academic Press, New York, 1980, pp. 675–715.
  90. Lightbrown, J. W. & Jackson, F. L., Inhibition of cytochrome systems of heart muscle and certain bacteria by the antagonists of dihydrostreptomycin: 2-alkyl-4-hydroxyquinoline *N*-oxides. *Biochem. J.*, **63** (1956) 130–7.
  91. Bowyer, J. R., Edwards, C. A., Ohnishi, T. & Trumpower, B. L., An analogue of ubiquinone which inhibits respiration by binding to the iron-sulfur protein of the cytochrome *bc*<sub>1</sub> segment of the mitochondrial respiratory chain. *J. Biol. Chem.*, **257** (1982) 8321–30.
  92. Oettmeier, W., Masson, K., Soll, M. & Reil, E., Acridones and quinolones as inhibitors of ubiquinone functions in the mitochondrial respiratory chain. *Biochem. Soc. Trans.*, **22** (1994) 213–6.
  93. Oettmeier, W., Dostatni, R., Majewski, C., Höfle, G., Fecker, T., Kunze, B. & Reichenbach, H., The aurachins, naturally occurring inhibitors of photosynthetic electron

flow through Photosystem II and the cytochrome  $b_6/f$ -complex. *Z. Naturforsch.*, **45** (1990) 322–8.

94. Arora, S. K., Bates, R. B., Grady, R. A. & Delfel, N. E.,

Crystal and molecular structure of the one to one complex of rotenone and carbon tetrachloride. *J. Am. Chem. Soc.*, **97** (1975) 5752–5.